

# onsior<sup>TM</sup> (robenacoxib)

10 mg, 20 mg & 40 mg  
Tablets for Dogs

For Oral Use In Dogs Only

**Information for Dog Owners** – You should read this information before starting your dog on ONSIOR tablets.

**Store ONSIOR out of reach of dogs and other pets in a secured location in order to prevent accidental ingestion or overdose.**

## What is ONSIOR?

ONSIOR (robenacoxib) is a prescription non-narcotic, non-steroidal anti-inflammatory (NSAID) of the coxib class. ONSIOR (robenacoxib) is for the control of postoperative pain and inflammation associated with soft tissue surgery (e.g. spay, neuter, tumor removal and stomach surgery) in dogs greater than  $\geq 5.5$  lbs (2.5 kg) and  $\geq 4$  months of age for a **maximum of 3 days**. **Do not use in dogs less than 5.5 lbs. Do not use in cats. Cats cannot be accurately dosed with the tablet sizes intended to be used in dogs.**

Your dog received the first dose of ONSIOR in the hospital prior to surgery. Your dog should not receive more than one dose a day, for a **maximum of 3 total doses**. Serious side effects have been associated with long term use. Follow the instructions given to you by your veterinarian.

# onsior<sup>TM</sup> (robenacoxib)

This summary contains important information about ONSIOR but does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or if you want to know more about ONSIOR for dogs.

## What kind of results can I expect when my dog takes ONSIOR for postoperative pain and inflammation?

ONSIOR may help your dog to recover more comfortably by controlling postoperative pain and inflammation following soft tissue surgery.

- Control of postoperative pain and inflammation may vary from dog to dog.
- Administer ONSIOR according to your veterinarian's directions, for a maximum of 3 days.
- Consult your veterinarian if your dog appears to be uncomfortable.

## Which dogs should not take ONSIOR?


Your dog should not be given ONSIOR if s/he:

- Has had an allergic reaction to robenacoxib, the active ingredient in ONSIOR
- Has had an allergic reaction (such as hives, facial swelling, or red or itchy skin) to aspirin or other NSAIDs
- Is less than 5.5 lbs (2.5 kg) and/or less than 4 months of age
- Has a loss of appetite/decreased appetite
- Has bloody feces or vomit
- Is presently taking aspirin, other NSAIDs, or corticosteroids
- Has a pre-existing liver or kidney condition
- Has any condition predisposing to dehydration

## ONSIOR Tablets for Dogs should only be given to dogs.

People should not take ONSIOR. Keep ONSIOR and all medication out of reach of children and pets. Call your physician immediately if you accidentally take ONSIOR.



 **Do not use ONSIOR dog tablets in cats. Cats cannot be accurately dosed with the ONSIOR dog tablet sizes intended to be used in dogs. ONSIOR tablets should not be broken.**

**Tell your veterinarian about:**

- Any side effects your dog has previously experienced from ONSIOR or other NSAIDs
- Any digestive upset (vomiting or diarrhea) your dog has had
- Any cardiovascular, kidney or liver disease your dog has had
- Any other medical problems or allergies that your dog has now or has had in the past
- All medications that you are giving your dog or plan to give your dog, including those you can get without prescription and any dietary supplements
- If you plan to breed your dog, or if your dog is pregnant or nursing

**Talk to your veterinarian about:**

- The surgery performed on your dog
- What test should be done before surgery and prior to administering ONSIOR
- The signs of postoperative pain or inflammation that may occur following surgery
- Normal events that can be expected after your dog undergoes surgery
- How often your dog may need to be examined by your veterinarian
- The risks and benefits of using ONSIOR.

**How to give ONSIOR to your dog.**

ONSIOR should be given according to your veterinarian's instructions. Your veterinarian will tell you what amount of ONSIOR is right for your dog. **ONSIOR should only be given once a day and for no longer than 3 days.** ONSIOR tablets should not be broken. ONSIOR tablets should be given by mouth and may be given with or without food.

**What are the possible side effects that may occur in my dog during therapy with ONSIOR?**

ONSIOR may cause some side effects in individual dogs. These are normally mild, but serious side effects have been reported in dogs taking non-steroidal anti-inflammatory drugs (NSAIDs) including ONSIOR. Serious side effects can result in death. It is important to stop the medication and contact your veterinarian immediately if you think your dog may have a medical problem or side effect while on ONSIOR. To report a possible side effect, contact your veterinarian or call Elanco US Inc. at 1-888-545-5973. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/AnimalVeterinary/SafetyHealth>.

Look for these possible side effects that may indicate that your dog is having a problem:

- Decrease in appetite
- Vomiting
- Change in bowel movements such as diarrhea or change in stool color
- Change in drinking or urination
- Change in behavior, such as depression or restlessness

**If you notice any of the above signs in your dog, stop administering ONSIOR and call your veterinarian.**

**Can ONSIOR be given with other medications?**

ONSIOR should not be given with other non-steroidal anti-inflammatory drugs (NSAIDs – e.g. aspirin, carprofen) or corticosteroids (e.g. prednisone).

Always tell your veterinarian about all medications that you have given your dog in the past, and any medications that you are planning to give with ONSIOR.

**What can I do in case my dog eats more than the prescribed amount of ONSIOR tablets?**

Contact your veterinarian immediately if your dog eats more than the prescribed amount of ONSIOR.

**What else should I know about ONSIOR?**

This sheet provides a summary of information about ONSIOR for Dogs. If you have any questions or concerns about ONSIOR or postoperative pain and inflammation, talk to your veterinarian.

As with all prescribed medications, ONSIOR should only be given to the dog for which it is prescribed. ONSIOR should be given to your dog only for the condition for which it was prescribed, at the prescribed dose, as directed by your veterinarian.

NADA 141-463, Approved by FDA

Elanco, ONSIOR and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

PA090267AMA 12267

**Elanco**